UG/5th Sem/Math(H)/T/19

2019

B.Sc. (Honours)

5th Semester Examination

MATHEMATICS

Paper - C11T

(Partial Differential Equations and Applications)

Full Marks: 60 Time: 3 Hours

The figures in the margin indicate full marks. Candidates are required to give their answers in their own words as far as practicable.

Unit - I

(1st Order PDE)

1. Answer any four questions:

- 2×4
- (a) Define Quasi-Linear PDE of first order and give an example.
- (b) Form a PDE by eliminating the arbitrary function φ from $Z = e^{ny} \varphi(x-y)$.

- of constant radius and having centre on the xy-plane.
 - (d) Find the characteristic curve of the PDE:

$$yz \frac{\partial z}{\partial x} + xz \frac{\partial z}{\partial y} = xy$$
,

- (e) Reduce the equation $u_x xu_y = 0$ in canonical form.
 - (f) Let z(x, y) be the solution of

 $x\frac{\partial z}{\partial x} + y\frac{\partial z}{\partial y} = 4z$ satisfying the condition z(x, y) = 1 on the circle $x^2 + y^2 = 1$. Then find the value of z(2, 2).

2. Answer any one question:

5×1

(a) Using the method of separation of variables solve $4\frac{\partial z}{\partial x} - \frac{\partial z}{\partial y} = 3z \text{ where } z(0, y) = 3e^{-y} - e^{-5y}.$

(b) Find the integral surface of the linear PDE:

$$x(y^2+z)\frac{\partial z}{\partial x} - y(x^2+z)\frac{\partial z}{\partial y} = (x^2-y^2)z$$

which contains the straight line x + y = 0, z = 1.

Unit - II

(2nd Order PDE)

3. Answer any one question:

 2×1

(a) Determine the region in which the given equation is hyperbolic, parabolic or elliptic

$$u_{xx} + xyu_{xyy} = 0.$$

(b) Find the characteristics of the PDE

$$\frac{\partial^2 z}{\partial x^2} - 2 \frac{\partial^2 z}{\partial x \partial y} + \frac{\partial^2 z}{\partial y^2} + \frac{\partial z}{\partial x} - \frac{\partial z}{\partial y} - e^x (2y - 3) + e^y = 0.$$

4. Answer any one question:

10×1

(a) (i) Reduce the equation

 $3u_{xx} + 10u_{xy} + 3u_{yy} = 0$ to its canonical form and hence solve it.

[Turn Over]

(ii) Derive the one dimensional wave equation.

6+4

(b) (i) Use the polar co-rodinates r and

 $\theta(x = r\cos\theta, y = r\sin\theta)$ to transform the Laplace equation $u_{xx} + u_{yy} = 0$ into the polar form

$$\nabla^2 u = u_{rr} + \frac{1}{r} u_r + \frac{1}{r^2} u_{\theta\theta} = 0$$

(ii) Reduce the Tricomi equation

 $u_{xx} + xu_{yy} = 0$ to the canonical form when x > 0.

Unit - III

(Applications of PDE)

5. Answer any one question:

5×1

(a) Find a solution of the following nonhomogeneous boundary value problem

$$u_{tt} = c^2 u_{xx} + F(x), \ 0 < x < l, \ t > 0$$

$$u(x, 0) = f(x), \quad 0 \le x \le l$$

$$u_t(x, 0) = g(x), 0 \le x \le l$$

 $u(0, t) = A, u(l, t) = B, t > 0,$ 5

(b) Find the solution of Laplace's equation $\nabla^2 \psi = 0$ in the semifinite region bounded by $x \ge 0$, $0 \le y \le 1$ subject to the boundary conditions

$$\left(\frac{\partial \Psi}{\partial x}\right)_{x=0} = 0, \ \left(\frac{\partial \Psi}{\partial y}\right)_{y=0} = 0 \ \text{and}$$

$$\psi(x, 1) = f(x).$$

6. Answer any one question:

10×1

(a) Obtain the D'Alembert's solution of the Cauchy problem

$$u_{tt} - c^2 u_{xx} = 0, \qquad x \in \mathbb{R}, \ t > 0$$

$$u(x, 0) = f(x), \quad x \in \mathbb{R}$$

$$u_t(x, 0) = g(x), \quad x \in \mathbb{R}$$

Define the term 'domain of dependence', 'range of influence'. Hence find the solution when

[Turn Over]

$$f(x) = |\sin x|, x > 0$$

= 0, x < 0
$$g(x) = 0, x \in \mathbb{R}$$

(b) Find the temperature distribution of a rod for the following initial boundary value problem

$$u_t = Ku_{xx}, \ 0 < x < l, \ t > 0$$
 $u(0, t) = 0, \ t \ge 0$
 $u(x, 0) = f(x), \ 0 \le x \le l$
 $u(l, t) = 0, \ t \ge 0$

Hence find the solution when f(x) = x(l-x), $0 \le x \le l$.

Unit - IV

(Particle Dynamics)

7. Answer any five questions:

 2×5

(a) Derive the relation $w = \frac{v \sin \varphi}{r} = \frac{vp}{r}$.

- (b) Prove that the acceleration of a particle moving in a plane curve with uniform speed is $\rho \psi^2$.
- (e) Write the Kepler's Law of planetary motion.
 - (d) For a particle moving in a central orbit under the inverse square law $\left(\frac{\mu}{r^2}\right)$, prove that the velocity (v) at any distance r is given by

$$v^2 = \mu \left(\frac{2}{r} - \frac{1}{a} \right).$$

- (e) Write the significance of $h = r^2 \dot{\theta}$.
- Write the differential equation of the central orbit in Pedal form.
- (g) A particle describes the parabola $p^2 = ar$ under a force, which is always directed towards its focus. Find the law of force.
 - (h) A point moves along the arc of a cycloid in such a manner that the tangent as it rotates with constant angular velocity. Show that the acceleration of the moving point is constant in magnitude.

[Turn Over]

8. Answer any two questions:

5×2

- (a) A particle describes an ellipse under a force which is always directed towards the centre of the ellipse. Find the law of force.
 - (b) A machine gun of mass M_o stands on a horizontal plane and contains a shot of total mass m which is fired horizontally at a uniform rate with constant velocity u relative to the gun.
 - (c) A particle is projected along the inner surface of a rough sphere and is acted on by no force. Show that it will return to the point of projection at the end of time $a/\mu V(e^{2\pi\mu}-1)$ where a is the radius of the sphere, V is the velocity of projection and μ is the coefficient of friction.